arXiv:1210.3875v1 [math.CA] 15 Oct 2012

Sharp Inequalities between Harmonic, Seiffert, Quadratic and Contraharmonic Means

Gen-Di Wang[1], Chen-Yan Yang[2] and Yu-Ming Chu[1]

[1]Department of Mathematics, Huzhou Teachers College, Huzhou 313000, China;
[2]Department of Mathematics, Zhejiang Sci-Tech University, Hangzhou 310018, China.
Correspondence should be addressed to Yu-Ming Chu, chuyuming@hutc.zj.cn

Abstract: In this paper, we present the greatest values α, λ and p, and the least values β, μ and q such that the double inequalities $\alpha D(a,b)+(1-\alpha)H(a,b)<T(a,b)<\beta D(a,b)+(1-\beta)H(a,b)$, $\lambda D(a,b)+(1-\lambda)H(a,b)<C(a,b)<\mu D(a,b)+(1-\mu)H(a,b)$ and $pD(a,b)+(1-p)H(a,b)<Q(a,b)<qD(a,b)+(1-q)H(a,b)$ hold for all $a,b>0$ with $a\neq b$, where $H(a,b)=2ab/(a+b)$, $T(a,b)=(a-b)/[2\arctan((a-b)/(a+b))]$, $Q(a,b)=\sqrt{(a^2+b^2)/2}$, $C(a,b)=(a^2+b^2)/(a+b)$ and $D(a,b)=(a^3+b^3)/(a^2+b^2)$ are the harmonic, Seiffert, quadratic, first contraharmonic and second contraharmonic means of a and b, respectively.

2010 Mathematics Subject Classification: 26E60.
Keywords: harmonic mean, Seiffert mean, quadratic mean, first contraharmonic mean, second contraharmonic mean.

1 Introduction

For $a,b>0$ with $a\neq b$ the harmonic mean $H(a,b)$, Seiffert mean $T(a,b)$, quadratic mean $Q(a,b)$, first contraharmonic mean $C(a,b)$ and second contraharmonic mean $D(a,b)$ are defined by $H(a,b)=2ab/(a+b)$, $T(a,b)=(a-b)/[2\arctan((a-b)/(a+b))]$, $Q(a,b)=\sqrt{(a^2+b^2)/2}$, $C(a,b)=(a^2+b^2)/(a+b)$ and $D(a,b)=(a^3+b^3)/(a^2+b^2)$, respectively. Recently, the harmonic, Seiffert, quadratic, first contraharmonic and second contraharmonic means have attracted the attention of many mathematicians. In particular, many remarkable inequalities for these means can be found in the literature [1-28]. Let $G(a,b)=\sqrt{ab}$, $L(a,b)=(a-b)/(\log a-\log b)$, $I(a,b)=1/e(b^b/a^a)^{1/(b-a)}$, $A(a,b)=(a+b)/2$, and $M_p(a,b)=[(a^p+b^p)/2]^{1/p}(p\neq 0)$ and $M_0(a,b)=\sqrt{ab}$ be the geometric, logarithmic, identric, arithmetic and p-th powers means of a and b, respectively. Then it is well known that the inequalities

$$\begin{aligned} H(a,b)=M_{-1}(a,b)<G(a,b)=M_0(a,b)<L(a,b)<I(a,b)\\ <A(a,b)=M_1(a,b)<T(a,b)<Q(a,b)=M_2(a,b)<C(a,b)<D(a,b)\end{aligned}$$

hold for all $a,b>0$ with $a\neq b$.

For all $a,b>0$ with $a\neq b$ Seiffert [29] established the double inequality $A(a,b)<T(a,b)<Q(a,b)$. Hästö [30] proved that the function $T(1,x)/M_p(1,x)$ is increasing in $[1,\infty)$ if $p\leq 1$. Chu et al. [31] gave the greatest values α and λ, and the least values β and μ such that the double inequalities $\alpha Q(a,b)+(1-\alpha)A(a,b)<T(a,b)<\beta Q(a,b)+(1-\beta)A(a,b)$ and $Q^{\lambda}(a,b)A^{1-\lambda}(a,b)<T(a,b)<Q^{\mu}(a,b)A^{1-\mu}(a,b)$ hold for all $a,b>0$ with $a\neq b$.

For $\alpha_1,\alpha_2,\beta_1,\beta_2\in(0,1/2)$, Chu et al. [32, 33] proved that the double inequalities

$$Q(\alpha_1a+(1-\alpha_1)b,\alpha_1b+(1-\alpha_1)a)<T(a,b)<Q(\beta_1a+(1-\beta_1)b,\beta_1b+(1-\beta_1)a)$$

and

$$C(\alpha_2a+(1-\alpha_2)b,\alpha_2b+(1-\alpha_2)a)<T(a,b)<C(\beta_2a+(1-\beta_2)b,\beta_2b+(1-\beta_2)a)$$

hold for all $a,b>0$ with $a\neq b$ if and only if $\alpha_1\leq(1+\sqrt{16/\pi^2-1})/2$, $\beta_1\geq(3+\sqrt{6})/6$, $\alpha_2\leq(1+\sqrt{4/\pi-1})/2$ and $\beta_2\geq(3+\sqrt{3})/6$.

In [34] Neuman proved that the double inequalities

$$\alpha Q(a,b)+(1-\alpha)A(a,b)<NS(a,b)<\beta Q(a,b)+(1-\beta)A(a,b)$$

and

$$\lambda C(a,b)+(1-\lambda)A(a,b)<NS(a,b)<\mu C(a,b)+(1-\mu)A(a,b)$$

hold for all $a,b>0$ with $a\neq b$ if and only if $\alpha\leq[1-\log(1+\sqrt{2})]/[(\sqrt{2}-1)\log(1+\sqrt{2})]=0.3249\cdots$, $\beta\geq 1/3$, $\lambda\leq[1-\log(1+\sqrt{2})]/\log(1+\sqrt{2})=0.1345\cdots$ and $\beta\geq 1/6$, where

$$NS(a,b)=\frac{a-b}{2\sinh^{-1}[(a-b)/(a+b)]}.$$

It is the aim of this paper to present the greatest values α, λ and p, and the least values β, μ and q such that the double inequalities

$$\alpha D(a,b)+(1-\alpha)H(a,b)<T(a,b)<\beta D(a,b)+(1-\beta)H(a,b)$$

$$\lambda D(a,b)+(1-\lambda)H(a,b)<C(a,b)<\mu D(a,b)+(1-\mu)H(a,b)$$

and

$$pD(a,b)+(1-p)H(a,b)<Q(a,b)<qD(a,b)+(1-q)H(a,b)$$

hold for all $a,b>0$ with $a\neq b$.

2 Lemmas

In order to prove our main results we need several lemmas, which we present in this section.

Lemma 2.1. Let $f_1(t)=(\pi^2-\pi-4)t^6-2\pi(5-\pi)t^5-3\pi(5-\pi)t^4-4(5\pi-\pi^2-2)t^3-3\pi(5-\pi)t^2-2\pi(5-\pi)t+(\pi^2-\pi-4)$. Then there exists $\lambda_0>1$ such that $f_1(t)<0$ for $t\in[1,\lambda_0)$ and $f_1(t)>0$ for $t\in(\lambda_0,+\infty)$.

Proof. Simple computations lead to

$$f_1(1)=-8\pi(9-2\pi)<0,\quad \lim_{t\to+\infty}f_1(t)=+\infty, \tag{2.1}$$

$$\begin{aligned}{f_1}'(t)=6(\pi^2-\pi-4)t^5-10\pi(5-\pi)t^4-12\pi(5-\pi)t^3\\ -12(5\pi-\pi^2-2)t^2-6\pi(5-\pi)t-2\pi(5-\pi),\end{aligned}$$

$${f_1}'(1)=-24\pi(9-2\pi)<0,\quad \lim_{t\to+\infty}{f_1}'(t)=+\infty, \tag{2.2}$$

$${f_1}''(t)=30(\pi^2-\pi-4)t^4-40\pi(5-\pi)t^3-36\pi(5-\pi)t^2-24(5\pi-\pi^2-2)t-6\pi(5-\pi).$$

$${f_1}''(1)=-8(70\pi+9-17\pi^2)<0,\quad \lim_{t\to\infty}{f_1}''(t)=+\infty, \tag{2.3}$$

$${f_1}'''(t)=120(\pi^2-\pi-4)t^3-120\pi(5-\pi)t^2-72\pi(5-\pi)t-24(5\pi-\pi^2-2),$$

$${f_1}'''(1)=-48(25\pi+9-7\pi^2)<0,\quad \lim_{t\to\infty}{f_1}'''(t)=+\infty, \tag{2.4}$$

$${f_1}^{(4)}(t)=360(\pi^2-\pi-4)t^2-240\pi(5-\pi)t-72\pi(5-\pi),$$

$$f_1{}^{(4)}(1) = -96(20\pi + 15 - 7\pi^2) < 0, \quad \lim_{t\to\infty} f_1{}^{(4)}(t) = +\infty, \tag{2.5}$$

$$\begin{aligned} f_1{}^{(5)}(t) =& 720(\pi^2 - \pi - 4)t - 240\pi(5-\pi) > 720(\pi^2 - \pi - 4) - 240\pi(5-\pi) \\ =& 960(\pi+1)(\pi-3) > 0 \end{aligned} \tag{2.6}$$

for $t > 1$.

Inequality (2.6) implies that $f_1{}^{(4)}(t)$ is strictly increasing in $[1, +\infty)$. Then from (2.5) we clearly see that there exists $\lambda_1 > 1$ such that $f_1{}^{(4)}(t) < 0$ for $t \in [1, \lambda_1)$ and $f_1{}^{(4)}(t) > 0$ for $t \in (\lambda_1, +\infty)$. Hence $f_1{}'''(t)$ is strictly decreasing in $[1, \lambda_1]$ and strictly increasing in $[\lambda_1, +\infty)$.

It follows from (2.4) and the piecewise monotonicity of $f_1{}'''(t)$ that there exists $\lambda_2 > \lambda_1 > 1$ such that $f_1{}''(t)$ is strictly decreasing in $[1, \lambda_2]$ and strictly increasing in $[\lambda_2, +\infty)$.

From (2.3) and the piecewise monotonicity of $f_1{}''(t)$ we clearly see that there exists $\lambda_3 > \lambda_2 > 1$ such that $f_1{}'(t)$ is strictly decreasing in $[1, \lambda_3]$ and strictly increasing in $[\lambda_3, +\infty)$.

The piecewise monotonicity of $f'(t)$ and (2.2) lead to the conclusion that there exists $\lambda_4 > \lambda_3 > 1$ such that $f_1(t)$ is strictly decreasing in $[1, \lambda_4]$ and strictly increasing in $[\lambda_4, +\infty)$.

Therefore, Lemma 2.1 follows from (2.1) and the piecewise monotonicity of $f_1(t)$.

Lemma 2.2. Let $c \in (0, 1)$, $t > 1$ and

$$f_c(t) = \frac{t^4 - 1}{ct^4 + (2-c)t^3 + (2-c)t + c} - 2\arctan\left(\frac{t-1}{t+1}\right). \tag{2.7}$$

Then $f_{4/9}(t) > 0$ and $f_{2/\pi}(t) < 0$ for all $t > 1$.

Proof. From (2.7) one has

$$f_c(1) = 0, \tag{2.8}$$

$$\lim_{t\to+\infty} f_c(t) = \frac{1}{c} - \frac{\pi}{2}, \tag{2.9}$$

$$f_c{}'(t) = \frac{g_c(t)}{(t^2+1)[ct^4 + (2-c)t^3 + (2-c)t + c]^2}, \tag{2.10}$$

where

$$\begin{aligned} g_c(t) =& (2 - 2c^2 - c)t^8 - 4c(2-c)t^7 + 2c(2-c)t^6 + 4c^2t^5 - 2(4c^2 - 5c + 2)t^4 \\ &+ 4c^2t^3 + 2c(2-c)t^2 - 4c(2-c)t + 2 - 2c^2 - c. \end{aligned} \tag{2.11}$$

We divide the proof into two cases.

Case 1 $c = 4/9$. Then (2.11) becomes

$$\begin{aligned} g_{4/9}(t) =& \frac{2}{81}(47t^8 - 112t^7 + 56t^6 + 32t^5 - 46t^4 + 32t^3 + 56t^2 - 112t + 47) \\ =& \frac{2}{81}(t-1)^4(47t^4 + 76t^3 + 78t^2 + 76t + 47) > 0 \end{aligned} \tag{2.12}$$

for $t > 1$.

Therefore, $f_{4/9}(t) > 0$ for $t > 1$ follows easily from (2.8) and (2.10) together with (2.12).

Case 2 $c = 2/\pi$. Then (2.9) and (2.11) lead to

$$\lim_{t\to+\infty} f_c(t) = \lim_{t\to+\infty} f_{2/\pi}(t) = 0, \tag{2.13}$$

$$\begin{aligned} g_{2/\pi}(t) =& \frac{2}{\pi^2}[(\pi^2 - \pi - 4)t^8 - 8(\pi - 1)t^7 + 4(\pi - 1)t^6 + 8t^5 - 2(\pi^2 - 5\pi + 8)t^4 \\ & + 8t^3 + 4(\pi - 1)t^2 - 8(\pi - 1)t + \pi^2 - \pi - 4] \\ =& \frac{2(t-1)^2}{\pi^2} f_1(t), \end{aligned} \tag{2.14}$$

where $f_1(t)$ is defined as in Lemma 2.1.

From (2.1) and (2.14) together with Lemma 2.1 we clearly see that $f_{2/\pi}(t)$ is strictly decreasing in $[1, \lambda_0]$ and strictly increasing in $[\lambda_0, +\infty)$.

Therefore, $f_{2/\pi}(t) < 0$ for $t > 1$ follows from (2.8) and (2.13) together with the piecewise monotonicity of $f_{2/\pi}(t)$.

Lemma 2.3. Let $t > 1$ and

$$g(t) = \frac{(t^2+1)[(t+1)\sqrt{t^2+1} - 2\sqrt{2}t]}{(t^2+t+1)(t-1)^2}. \tag{2.15}$$

Then $g(t)$ is strictly increasing from $(1, \infty)$ onto $(\sqrt{2}/2, 1)$.

Proof. From (2.15) we get

$$\lim_{t\to1^+} g(t) = \frac{\sqrt{2}}{2}, \quad \lim_{t\to+\infty} g(t) = 1. \tag{2.16}$$

Let $t = \tan x$, $u = \sin x + \cos x$. Then $x \in (\pi/4, \pi/2)$, $u \in (1, \sqrt{2})$ and (2.15) becomes

$$\begin{aligned} g(t) =& \frac{\sin x + \cos x - \sqrt{2}\sin(2x)}{(1+\sin(2x)/2)(1-\sin(2x))} \\ =& \frac{\sin x + \cos x - \sqrt{2}[(\sin x + \cos x)^2 - 1]}{\{1 + [(\sin x + \cos x)^2 - 1]/2\}[2 - (\sin x + \cos x)^2]} \\ =& \frac{2(u - \sqrt{2}u^2 + \sqrt{2})}{-u^4 + u^2 + 2} \triangleq h(u). \end{aligned} \tag{2.17}$$

Equation (2.17) leads to

$$\begin{aligned} h'(u) =& \frac{-4\sqrt{2}u^5 + 6u^4 + 8\sqrt{2}u^3 - 2u^2 - 12\sqrt{2}u + 4}{(-u^4 + u^2 + 2)^2} \\ =& \frac{-4\sqrt{2}(u - \sqrt{2})^2}{(-u^4 + u^2 + 2)^2}\left(u^3 + \frac{5\sqrt{2}}{4}u^2 + u - \frac{\sqrt{2}}{4}\right) < 0. \end{aligned} \tag{2.18}$$

Note that $x \to \sin x + \cos x$ is strictly decreasing from $(\pi/4, \pi/2)$ onto $(1, \sqrt{2})$. Therefore, Lemma 2.3 follows easily from (2.16)-(2.18).

3 Main Results

Theorem 3.1. The double inequality

$$\alpha D(a,b) + (1-\alpha)H(a,b) < T(a,b) < \beta D(a,b) + (1-\beta)H(a,b) \tag{3.1}$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha \leq 4/9$ and $\beta \geq 2/\pi$.

Proof. Since $H(a,b)$, $T(a,b)$ and $D(a,b)$ are symmetric and homogeneous of degree 1. Without loss of generality, we assume that $a > b$. Let $t = a/b > 1$ and $c \in (0,1)$. Then simple computations lead to

$$\frac{T(a,b) - H(a,b)}{D(a,b) - H(a,b)} = \frac{(t^2+1)\left(t^2 - 4t\arctan\left(\frac{t-1}{t+1}\right) - 1\right)}{2(t-1)^2(t^2+t+1)\arctan\left(\frac{t-1}{t+1}\right)}, \tag{3.2}$$

$$\lim_{t\to 1^+} \frac{(t^2+1)\left(t^2 - 4t\arctan\left(\frac{t-1}{t+1}\right) - 1\right)}{2(t-1)^2(t^2+t+1)\arctan\left(\frac{t-1}{t+1}\right)} = \frac{4}{9}, \tag{3.3}$$

$$\lim_{t\to+\infty}\frac{(t^2+1)\left(t^2-4t\arctan\left(\frac{t-1}{t+1}\right)-1\right)}{2(t-1)^2(t^2+t+1)\arctan\left(\frac{t-1}{t+1}\right)}=\frac{2}{\pi}, \tag{3.4}$$

$$\begin{aligned}&T(a,b)-[cD(a,b)+(1-c)H(a,b)]\\&\quad=b\frac{c(t+1)(t^3+1)+2(1-c)t(t^2+1)}{2(t+1)(t^2+1)\arctan\left(\frac{t-1}{t+1}\right)}f_c(t),\end{aligned} \tag{3.5}$$

where $f_c(t)$ is defined as in Lemma 2.2.

Therefore, inequality $4D(a,b)/9+5H(a,b)/9<T(a,b)<2D(a,b)/\pi+(1-2/\pi)H(a,b)$ holds for all $a,b>0$ with $a\neq b$ follows from (3.5) and Lemma 2.2.

Next, we prove that $\alpha=4/9$ and $\beta=2/\pi$ are the best possible parameters such that inequality (3.1) holds for all $a,b>0$ with $a\neq b$.

If $\alpha>4/9$, then from (3.2) and (3.3) we know that there exists $\delta>0$ such that $T(a,b)<\alpha D(a,b)+(1-\alpha)H(a,b)$ for all $a,b>0$ with $a/b\in(1,1+\delta)$.

If $\beta<2/\pi$, then (3.2) and (3.3) lead to the conclusion that there exists $T_0>1$ such that $T(a,b)>\beta D(a,b)+(1-\beta)H(a,b)$ for all $a,b>0$ with $a/b\in(T_0,\infty)$.

Theorem 3.2. The double inequality

$$pD(a,b)+(1-p)H(a,b)<Q(a,b)<qD(a,b)+(1-q)H(a,b)$$

holds for all $a,b>0$ with $a\neq b$ if and only if $p\leq 1/2$ and $q\geq\sqrt{2}/2$.

Proof. Since $H(a,b)$, $Q(a,b)$ and $D(a,b)$ are symmetric and homogeneous of degree 1. Without loss of generality, we assume that $a>b$. Let $t=a/b>1$, then

$$\frac{Q(a,b)-H(a,b)}{D(a,b)-H(a,b)}=\frac{(t^2+1)[(t+1)\sqrt{t^2+1}-2\sqrt{2}t]}{\sqrt{2}(t^2+t+1)(t-1)^2}=\frac{\sqrt{2}}{2}g(t), \tag{3.6}$$

where $g(t)$ is defined as in Lemma 2.3.

Therefore, Theorem 3.2 follows directly from (3.6) and Lemma 2.3.

Theorem 3.3. The double inequality

$$\lambda D(a,b)+(1-\lambda)H(a,b)<C(a,b)<\mu D(a,b)+(1-\mu)H(a,b)$$

holds for all $a,b>0$ with $a\neq b$ if and only if $\lambda\leq 2/3$ and $\mu\geq 1$.

Proof. Since $H(a,b)$, $C(a,b)$ and $D(a,b)$ are symmetric and homogeneous of degree 1. Without loss of generality, we assume that $a > b$. Let $t = a/b > 1$, then simple computations lead to

$$\frac{C(a,b) - H(a,b)}{D(a,b) - H(a,b)} = \frac{t^2+1}{t^2+t+1}, \tag{3.7}$$

$$\lim_{t\to 1^+} \frac{t^2+1}{t^2+t+1} = \frac{2}{3}. \tag{3.8}$$

$$\lim_{t\to +\infty} \frac{t^2+1}{t^2+t+1} = 1. \tag{3.9}$$

Note that the function $t \to (t^2+1)/(t^2+t+1)$ is strictly increasing in $[1, \infty)$. Therefore, Theorem 3.3 follows from (3.7)-(3.9) and the monotonicity of the function $t \to (t^2+1)/(t^2+t+1)$.

Acknowledgements. This research was supported by the Natural Science Foundation of China under Grants 11071069 and 11171307, and the Innovation Team Foundation of the Department of Education of Zhejiang Province under Grant T200924.

References

[1] W. Gautschi, An harmonic mean inequality for the gamma function, SIAM J. Math. Anal., 1974, **5**: 278-281.

[2] H. Alzer, Inequalities for arithmetic, geometric and harmonic means, Bull. London Math. Soc., 1990, **22**(4): 362-366.

[3] H. Alzer, An inequality for arithmetic and harmonic means, Aequationes Math., 1993, **46**(3): 257-263.

[4] H. Alzer, A harmonic mean inequality for the gamma function, J. Comput. Appl. Math., 1997, **87**(2): 195-198.

[5] Gh. Toader, Seiffert type means, Nieuw Arch. Wisk. (4), 1999, **17**(3): 379-382.

[6] H. Alzer, On Gautsch's harmonic mean inequality for the gamma function, J. Comput. Appl. Math., 2003, **157**(1): 243-249.

[7] P. R. Mercer, Refined arithmetic, geometric and harmonic mean inequalities, Rocky Mountain J. Math., 2003, **33**(4): 1459-1464.

[8] R. Sharma, Some inequalities for arithmetic mean, harmonic mean and variance, J. Math. Inequal., 2008, **2**(1): 109-114.

[9] Y.-M. Chu and W.-F. Xia, Two sharp inequalities for power mean, geometric mean, and harmonic mean, J. Inequal. Appl., 2009, Article ID 741923, 6 pages.

[10] Y.-M. Chu and W. F. Xia, Inequalities for generalized logarithmic means, J. Inequal. Appl., 2009, Article ID 763252, 7 pages.

[11] Y.-M. Chu, Y.-F. Qiu, M.-K. Wang and G.-D. Wang, The optimal convex combination bounds of arithmetic and harmonic means for the Seiffert's mean, J. Inequal. Appl., 2010, Article ID 436457, 7 pages.

[12] Y.-M. Chu, M.-K. Wang and Y.-F. Qiu, An optimal double inequalities between power-type Heron and Seiffert means, J. Inequal. Appl., 2010, Article ID 146945, 11 pages.

[13] B.-Y. Long and Y.-M. Chu, Optimal power mean bounds for the weighted geometric mean of classical means, J. Inequal. Appl., 2010, Article ID 905679, 6 pages.

[14] Y.-M. Chu and B.-Y. Long, Best possible inequalities between generalized logarithmic mean and classical means, Abstr. Appl. Anal., 2010, Article ID 303286, 13 pages.

[15] Y.-M. Chu, M.-K. Wang, S.-L. Qiu and Y.-F. Qiu, Sharp generalized Seiffert mean bounds for Toader mean, Abstr. Appl. Anal., 2011, Article ID 605259, 8 pages.

[16] Y.-M. Chu, M.-K. Wang and Z.-K. Wang, A sharp double inequality between harmonic and identric means, Abstr. Appl. Anal., 2011, Article ID 657935, 7 pages.

[17] Y.-M. Chu and M.-K. Wang, Optimal inequalities between harmonic, geometric, logarithmic, and arithmetic-geometric means, J. Appl. Math., 2011, Article ID 618929, 9 pages.

[18] Y.-M. Chu, M.-K. Wang and Z.-K. Wang, A best-possible double inequality between Seiffert and harmonic means, J. Inequal. Appl., 2011, **2011**:94, 7 pages.

[19] Y.-M. Chu, S.-W. Hou and W.-M. Gong, Inequalities between logarithmic, harmonic, arithmetic and centroidal means, J. Math. Anal., 2011, **2**(2): 1-5.

[20] Y.-M. Chu and B.-Y. Long, Sharp inequalities between means, Math. Inequal. Appl., 2011, **14**(3): 647-655.

[21] Y.-M. Chu, S.-S. Wang and C. Zong, Optimal lower power mean bounds for the convex combination of harmonic and logarithmic means, Abstr. Appl. Anal., 2011, Article ID 520648, 9 pages.

[22] S.-Q. Gao, Inequalities for the Seiffert's means in terms of the identric means, J. Math. Sci. Adv. Appl., 2011, **10**(1-2): 23-31.

[23] H. Liu and X.-J. Meng, The optimal convex combination bounds for the Seiffert's mean, J. Inequal. Appl., 2011, Article ID 686834, 9 pages.

[24] Y.-M. Chu, C. Zong and G.-D. Wang, Optimal convex combination bounds of Seiffert and geometric means for the arithmetic mean, J. Math. Inequal., 2011, **5**(3): 429-434.

[25] W.-F. Xia, W. Janous and Y.-M. Chu, The optimal convex combination bounds of arithmetic and harmonic means in terms of power mean, J. Math. Inequal., 2012, **6**(2): 241-248.

[26] Y.-M. Chu, M.-K. Wang and S.-L. Qiu, Optimal combinations bounds of root-square and arithmeitc means for Toader mean, Proc. Indian Acad. Sci. Math. Sci., 2012, **122**(1): 41-51.

[27] Y.-M. Chu, M.-K. Wang and Z.-K. Wang, Best possible inequalities among harmonic, geometric, logarithmic and Seiffert means, Math. Inequal. Appl., 2012, **15**(2): 415-422.

[28] Y.-M. Chu, M.-K. Wang and G.-D. Wang, The optimal generalized logarithmic mean bounds for Seiffert's mean, Acta Math. Sci., 2012, **32B**(4): 1619-1626.

[29] H.-J. Seiffert, Aufgabe β 16, Die Wurzel, 1995, **29**: 221-222.

[30] P. A. Hästö, A monotonicity property of ratios of symmetric homogeneous means, JIPAM. J. Inequal. Pure Appl. Math., 2002, **3**(5), Article 71, 23 pages.

[31] Y.-M. Chu, M.-K. Wang and W.-M. Gong, Two sharp double inequalities for Seiffert mean, J. Inequal. Appl., 2011, **2011**:44, 7 pages.

[32] Y.-M. Chu, S.-W. Hou and Z.-H. Shen, Sharp bounds for Seiffert mean in terms of root mean square, J. Inequal. Appl., 2012, **2012**:11, 6 pages.

[33] Y.-M. Chu and S.-W. Hou, Sharp bounds for Seiffert mean in terms of contraharmonic mean, Abstr. Appl. Anal., 2012, Article ID 425175, 6 pages.

[34] E. Neuman, A note on a certain bivariate mean, J. Math. Inequal., 2012, **6**(4): 637-643.